



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Physics, Computer Science & Mathematics Division

RECEIVED
LAWRENCE

BERKELEY LABORATORY

AUG 01 1980

LIBRARY AND
DOCUMENTS SECTION

Submitted for publication

LIMIT ON π MUOPRODUCTION AT 209 GeV

A.R. Clark, K.J. Johnson, L.T. Kerth, S.C. Loken,
T.W. Markiewicz, P.D. Meyers, W.H. Smith, M. Strovink,
W.A. Wenzel, R.P. Johnson, C. Moore, M. Mugge,
R.E. Shafer, G.D. Gollin, F.C. Shoemaker, and P. Surko

June 1980

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Limit on T Muoproduction at 209 GeV

A.R. Clark, K.J. Johnson, L.T. Kerth, S.C. Loken, T.W. Markiewicz,

P.D. Meyers, W.H. Smith, M. Strovink, and W.A. Wenzel

Physics Department and Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720

and

R.P. Johnson, C. Moore, M. Mugge, and R.E. Shafer

Fermi National Accelerator Laboratory,
Batavia, Illinois 60510

and

G.D. Gollin^a, F.C. Shoemaker, and P. Surko^b

Joseph Henry Laboratories, Princeton University,
Princeton, New Jersey 08544

We present the dimuon mass spectrum from 102 678 three-muon final states produced by muon interactions within a magnetized steel calorimeter. The data place a 90%-confidence limit on the production of T states by muons: $\sigma(\mu N \rightarrow \mu T X) B(T \rightarrow \mu^+ \mu^-) < 22 \times 10^{-39} \text{ cm}^2$, consistent with a photon-gluon-fusion model calculation.

We report a limit on T production by 209-GeV muons in the Berkeley-Fermilab-Princeton MultimMuon Spectrometer at Fermilab¹. An integrated luminosity of $0.78 \times 10^{39} \text{ cm}^{-2}$, corresponding to 75% of the full data sample, has yielded 102 678 trimuon final states, including 6693 ± 355 examples of J/ψ and ψ' production. In every event, all three outgoing muons are fully momentum-analyzed and are subjected to an energy-conserving one-constraint fit using calorimetric measurement of the associated shower energy.

No limit on T production by real or virtual photons has been published. A conference report² based on results from the Bologna-CERN-Dubna-Munich-Saclay (BCDMS) experiment presents the limit $\sigma(\mu N \rightarrow TX) B(T \rightarrow \mu^+ \mu^-) < (6 \pm 3) \times 10^{-39} \text{ cm}^2$ (90% confidence) for ~ 275 -GeV muons, where the error is systematic. This limit is based on 761 multimMuon events corresponding to an integrated luminosity² of $0.7 \times 10^{39} \text{ cm}^{-2}$. A third muon was observed in 11% of these events. No calorimetric information was available. With 48% T acceptance, the BCDMS limit corresponds to ≤ 2 T candidates (90% confidence). In total, the experiment observed 24 events between 8 and 12 GeV/c^2 in dimuon mass. These were compared to a calculated background of 30 electromagnetic triplets in the same region.

We have calculated the expected T rates using a photon-gluon-fusion (γGF) model³ which accounts⁴ for most of the published features¹ of ψ muoproduction. It uses a Bethe-Heitler diagram for heavy quark pair production with the nuclear photon replaced by a gluon. Additional soft gluon exchanges needed to conserve color are assumed not to affect the kinematics. Using a distribution $G(x) = 3(1-x)^5/x$ in gluon momentum fraction x , a bottom quark mass $m_b = 4.7 \text{ GeV}/c^2$, a bottom quark charge $|q_b| = 1/3$, and a strong coupling

constant $\alpha_s = 1.5 / \ln(4m_{b\bar{b}}^2)$, where $m_{b\bar{b}}$ is the mass in GeV/c^2 of the produced quark pair, the model predicts T muoproduction cross sections of $0.13 \times 10^{-36} \text{ cm}^2$ at 209 GeV and $0.28 \times 10^{-36} \text{ cm}^2$ at 275 GeV. With $B(T \rightarrow \mu^+ \mu^-) = (3.1 \pm 0.9)\%$ ⁵, the expected values of $B\sigma$ are $(4.0 \pm 1.2) \times 10^{-39}$ and $(8.7 \pm 2.5) \times 10^{-39} \text{ cm}^2$, respectively. The BCDMS upper limit is $(70 \pm 40)\%$ of the latter cross section.

Figure 1 displays the spectrum in dimuon mass $M_{\mu^+\mu^-}$ from this experiment. Events below $5 \text{ GeV}/c^2$ in $M_{\mu^+\mu^-}$ are reconstructed and momentum fit as described in Ref. 1. Above $5 \text{ GeV}/c^2$, the analysis of all events was checked by a hand reconstruction which was blind to the invariant mass. At all masses the assignment of beam-sign secondary muons either to the scattered muon or to the produced muon pair is the critical decision in the analysis. Incorrect pairing of muons from ψ or muon trident production can cause events which properly belong in the low-mass region to be misinterpreted as having a higher mass. Our muon pairing algorithm was selected primarily to minimize this problem. The scattered muon is chosen to be the one with the smaller value of the square of its scattering angle divided by its scattered energy. The algorithm is 89% efficient in reconstructing T 's generated by the Monte Carlo simulation described below. The alternative choice for the scattered muon would produce more than a one-order-of-magnitude exaggeration of the high-mass continuum near the T , as shown by the "mispaired" histogram segment in Fig. 1. We emphasize that the muon pairing algorithm can be optimized only if all three final-state muons are momentum-analyzed.

Despite the care exercised in muon pairing, Monte Carlo studies show that there remains a significant contribution in the region $4.7 < M_{\mu^+\mu^-} < 8.4 \text{ GeV}/c^2$

from incorrectly analyzed lower-mass events. Allowance for these effects is most reliably made by use of an empirical fit to the mass continuum. This mass region, together with the range $1.5 < M_{\mu^+\mu^-} < 2.3 \text{ GeV}/c^2$, was chosen for the fit in order to exclude regions complicated by charmonium production or rapid variations in low-mass acceptance. After subtraction of the fit continuum, the ψ peak in Fig. 1 exhibits an 8.5% rms resolution, $\approx 1\%$ larger than the Monte Carlo prediction¹. The extrapolated continuum contains 1.8 ± 1.0 background events in the T region $8.4 < M_{\mu^+\mu^-} < 11.1 \text{ GeV}/c^2$, which in fact includes two observed events. The additional event at $11.5 \text{ GeV}/c^2$ is interpreted as continuum background with 65% probability, or as part of the peak corresponding to known T states with 1% probability. With 90% confidence, there are fewer than 3.8 events above the extrapolated background.

The Monte Carlo program used to simulate T muoproduction is based on a routine which successfully parameterizes our ψ data¹. It is adapted to T simulation by appropriately scaling the vector-meson-mass-dependent parameters. Simulated T mass resolution and detection efficiency are 9% (rms) and 22%, respectively. The T cross section is normalized to the γGF value described above. T, T', and T'' states are generated in the ratio 1:0.39:0.32 in agreement with recent measurements of $\Gamma_{ee}(\text{T}):\Gamma_{ee}(\text{T}'):\Gamma_{ee}(\text{T}'')$ ⁶. T' and T'' production suffer an additional $\approx 30\%$ suppression relative to T production because of threshold kinematics. The reconstructed peak corresponding to $10^4 \times$ the expected signal is shown in Fig. 1; 1.0 events from all T states are expected in the data.

Our 3.8-event limit, integrated luminosity, and detection efficiency combine to set the 90%-confidence limit⁷ $\sigma(\mu\text{N} \rightarrow \mu\text{TX})\text{B}(\text{T} \rightarrow \mu^+\mu^-) < 22 \times 10^{-39} \text{ cm}^2$. With

$B(T \rightarrow \mu^+ \mu^-) = (3.1 \pm 0.9)\%$ ⁵, we obtain the 90%-confidence cross-section limit $\sigma(\mu N \rightarrow \mu TX) < 0.79 \times 10^{-36} \text{ cm}^2$, including the error in the branching ratio. This limit lies above published predictions which use either the vector-meson dominance^{8,9} or the γ GF¹⁰ models. Ignoring any γ GF model uncertainty, this result rules out the choice $|q_b| = 2/3$ with 85% confidence. With 67% confidence, the data disfavor the existence of similar bound states of a second charge 1/3 quark in the T mass region.

We are grateful for the dedicated efforts of many individuals at our respective institutions, and in the Fermilab Neutrino Department. This work was supported by the High Energy Physics Division of the U.S. Department of Energy under Contract Nos. W-7504-Eng-48, DE-AC02-76ER03072, and EY-76-C-02-3000.

References

- ^aNow at Enrico Fermi Institute, Chicago, IL 60637.
- ^bNow at Bell Laboratories, Murray Hill, NJ 07974.
- ¹A.R. Clark et al., Phys. Rev. Lett. 43, 187 (1979).
- ²A. Benvenuti, in Proc. Int. Symp. on Lepton and Photon Interactions at High Energies, edited by T.B.W. Kirk and H. Abarbanel (Fermilab, Batavia, Illinois, 1979), p. 149. The integrated luminosity is assumed to be the product $(1.5 \times 10^{11} \text{ muons}) \times (10 \text{ modules}) \times (500 \text{ cm of carbon/module}) \times (\text{density } 1.55 \text{ g/cm}^3) \times 6 \times 10^2$.
- ³J.P. Leveille and T. Weiler, Nucl. Phys. B147, 147 (1979), and references cited therein. In this model the fraction of bound heavy quarks in the 1S state is perhaps best regarded as a fit parameter. In agreement with ψ data (Ref. 1) we use the value 1/6. See V. Barger, W.Y. Keung, and R.J.N. Phillips, Univ. of Wisconsin Report No. 79-0776 (unpublished).
- ⁴T. Weiler, Phys. Rev. Lett. 44, 304 (1980).
- ⁵K. Berkelman, summary of DORIS results presented to the XV Rencontre de Moriond (Les Arcs, Savoie, France, March 9-21, 1980). If the bottom quark were to have charge 2/3, a substantially larger branching ratio would be expected.
- ⁶T. Bohringer et al., Phys. Rev. Lett. 44, 111 (1980). Due to lack of information we simply assume that T' and T'' have the $T \rightarrow \mu^+ \mu^-$ branching ratio.
- ⁷Strictly speaking, our limit is on $\sum_V \sigma(V) B(V \rightarrow \mu^+ \mu^-)$, where V runs over T , T' , and T'' . Our T' and T'' assumptions would then lead to a limit on T production which is 36% stronger than the limit we have quoted.
- ⁸G. Aubrecht and W. Wada, Phys. Rev. Lett. 39, 978 (1977), make photoproduc-

tion predictions which, with γ_{GF} ν - and Q^2 -dependence assumptions, correspond to 0.15×10^{-36} cm² of T muoproduction at 209 GeV.

⁹N. Bralić, Nucl. Phys. B139, 433 (1978), makes leptoproduction predictions which, with γ_{GF} energy-dependence assumptions, correspond to 0.07×10^{-36} cm² of T muoproduction at 209 GeV.

¹⁰H. Fritzsch and K. Streng, Phys. Lett. 72B, 385 (1978), make photoproduction predictions which, with γ_{GF} ν - and Q^2 -dependence assumptions, correspond to 0.36×10^{-36} cm² of T muoproduction at 209 GeV.

Figure Caption

FIG. 1. Spectrum of 102 678 dimuon masses from 75% of the trimuon data. The background is fit by $\exp(\alpha+bM+cM^2)$ in the regions of the solid curve with a χ^2 of 13.7 for 14 degrees of freedom, and is extrapolated along the dotted curve. The "mispaired" histogram segment illustrates the appearance of the mass spectrum if the alternative muon-pairing choice is made. The background-subtracted ψ peak is shown in the lower corner; the expected peak from $10^4 \times$ the Monte-Carlo simulated T , T' , and T'' sample is shown in the upper corner, with the contribution from T' and T'' in black.




FIG. 1

XBL 805-966